



Tree Survey

At

**Pontyfelin Lane,
New Inn,
Pontypool**

*Inspected by:-
Julian Wilkes BSc.For, MSc.Land Man, MIC.For, TechArborA
Treescene Ltd
The Walled Garden
Old Coedarhydyglyn
St Nicholas
Cardiff
CF5 6SG
Tel No. 029 20599300*

13th October, 2022

**Registered Office: Treescene Limited
The Walled Garden, Old Coedarhydyglyn, St. Nicholas, Cardiff CF5 6SG
Tel. 029 205 99300 Email. trees@treescene.co.uk**

Brief

I have been instructed by Mr Owain Morgan of Arup to carry out a survey on trees at Pontyfelin Lane, New Inn, Pontypool.

Scope of Report

This Tree Survey has been undertaken within the recommendations of British Standards 5837:2012 and current good arboricultural practice.

The survey entailed a visual inspection from ground level of all trees.

Each tree has been numbered and, where instructed, have been tagged using small durable metal or plastic tags.

Due to variations of existing ground levels through the site, height dimensions are estimated and are given in metres.

Trunk/stem diameters are measured at 1.5 metres above ground level, or immediately above the root flare for multi-stemmed trees.

Estimated branch spread is taken in metres from the centre of the trunk, at the four cardinal points of a compass, to achieve an accurate representation of crown shape.

An assessment of a tree's age classification is made in terms of its maturity within the site's landscape.

An assessment of a tree's physiological condition is made as good, fair, poor, dead.

Data on the structural condition of the tree has been entered, e.g., collapsing, leaning and the presence of any decay or physical defect has been noted.

Preliminary management recommendations include further investigation of suspected defects that require more detailed assessment or potential for wildlife habitat.

An assessment of a tree's future life expectancy is made as <10, 10-20, 20-40 or >40 etc.

Table 1 – Cascade chart for tree quality assessment

Category and definition	Criteria (including subcategories where appropriate)		
<p><u>Category U</u> Those in such a condition that they cannot realistically be retained as living trees in the context of the current land use for longer than 10 years</p>	<ul style="list-style-type: none"> • Trees that have a serious, irremediable, structural defect, such that their early loss is expected due to collapse, including those that will become unviable after removal of other U category trees (i.e. where, for whatever reason, the loss of companion shelter cannot be mitigated by pruning) • Trees that are dead or are showing signs of significant, immediate, and irreversible overall decline • Trees infected with pathogens of significance to the health and/or safety of other trees nearby, or very low quality trees suppressing adjacent trees of better quality <p>NOTE Category U trees can have existing or potential conservation value which it might be desirable to preserve; see 4.5.7</p>		
	1 Mainly Arboricultural values	2 Mainly landscape values	3 Mainly cultural values, including conservation
<p><u>Category A</u> Those of high quality with an estimated remaining life expectancy of at least 40 years</p>	Trees that are particularly good examples of their species, especially if rare or unusual, or essential components of groups, or of formal or semi-formal arboricultural features (e.g. the dominant and/or principal trees within an avenue)	Trees, groups or woodlands of particular visual importance as Arboricultural and/or landscape features	Trees, groups or woodlands of significant conservation; historical, commemorative or other value (e.g. veteran trees or wood-pasture)
<p><u>Category B</u> Those of moderate quality with an estimated remaining life expectancy of at least 20 years</p>	Trees that might be included in category A, but are downgraded because of impaired condition (e.g. presence of significant though remediable defects, including unsympathetic past management and storm damage), such that they are unlikely to be suitable for retention for beyond 40 years; or trees lacking the special quality necessary to merit the category A designation	Trees present in numbers, usually growing as groups or woodlands, such that they attract a higher collective rating than they might as individuals; or trees occurring as collectives but situated so as to make little visual contribution to the wider locality	Trees with material conservation or other cultural benefits
<p><u>Category C</u> Those of low quality with an estimated remaining life expectancy of at least 10 years, or young trees with a stem diameter below 150mm</p>	Unremarkable trees of very limited merit or such impaired condition that they do not qualify in higher categories	Trees present in groups or woodlands, but without this conferring on them significantly greater collective landscape value, and/or trees offering low or only temporary/transient landscape benefits	Trees with no material conservation or other cultural value

BRITISH STANDARD BS 5837:2012

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G1	Group of: Hawthorn (<i>Crataegus monogyna</i>), Hazel (<i>Corylus avellana</i>)	9	Single and Multi	0.25 (avg.)	5	2	3	3	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow with crowns more heavily developed on northern side.	No action required at this time.	20-40	C
T2	Ash (<i>Fraxinus excelsior</i>)	13	Multi	0.5	5	3	2	5	4	Middle aged	Poor	Field boundary specimen exhibiting early stage symptoms of Ash Dieback Disease.	Remove.	<10	U
T3	Hawthorn (<i>Crataegus monogyna</i>)	7	Single	0.19	1	1	1	4	2	Middle aged	Fair to poor	Scrubby field boundary specimen leaning to the west.	Monitor for stability.	10-20	C
G4	Group of: Ash (<i>Fraxinus excelsior</i>)	14	Multi	0.4	6	3	3	3	1	Middle aged	Poor	Field boundary trees infected with Ash Dieback Disease.	Remove.	<10	U
T5	Sycamore (<i>Acer pseudoplatanus</i>)	8	Single	0.17	3	1	1	3	2	Young	Fair	Scrubby field boundary specimens of variable form.	No action required at this time.	20-40	C
G6	Group of: Ash (<i>Fraxinus excelsior</i>)	20	Multi	0.75 (avg.)	11	8	8	10	3	Mature	Poor	Notable field boundary specimens exhibiting symptoms of Ash Dieback Disease.	Remove.	<10	U
G7	Group of: Hawthorn (<i>Crataegus monogyna</i>), Hazel (<i>Corylus avellana</i>), Holly (<i>Ilex aquifolium</i>)	8	Multi	0.2	2	2	2	2	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	20-40	C

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G8	Group of: Hazel (<i>Corylus avellana</i>), Hawthorn (<i>Crataegus monogyna</i>), Cherry (<i>Prunus</i> spp.)	3	Single and Multi	0.1	1	1	1	1	0	Young	Fair	Scrubby remnant section of hedgerow.	No action required at this time.	10-20	C
T9	Sycamore (<i>Acer pseudoplatanus</i>)	17	Single	0.44	3	7	6	7	3	Middle aged	Poor	Hedgerow tree of variable form exhibiting significant basal decay and internal decay associated with poor quality pruning wound on main stem at 1.5 metres. This specimen is potentially unstable.	Remove.	<10	U
T10	Ash (<i>Fraxinus excelsior</i>)	12	Single	0.14	1	3	2	1	2	Young	Poor	Hedgerow tree severely infected with Ash Dieback Disease.	Remove.	<10	U
T11	Crack Willow (<i>Salix fragilis</i>)	19	Multi	0.7	0	12	10	2	1	Mature	Poor	Hedgerow tree that is partially collapsed. This specimen is vulnerable to immediate failure.	Remove.	<10	U
G12	Group of: Hazel (<i>Corylus avellana</i>), Hawthorn (<i>Crataegus monogyna</i>)	4	Single and Multi	0.15	1	1	2	1	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	20-40	C
T13	Ash (<i>Fraxinus excelsior</i>)	20	Multi	0.55	9	6	6	7	4	Middle aged	Poor	Prominent field boundary hedgerow infected with Ash Dieback Disease.	Remove.	<10	U

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G14	Group of: Field Maple (<i>Acer campestre</i>), Holly (<i>Ilex aquifolium</i>), Hazel (<i>Corylus avellana</i>), Hawthorn (<i>Crataegus monogyna</i>)	10	Single and Multi	0.2 (avg.)	4	3	4	3	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	20-40	C
G15	Group of: Hawthorn (<i>Crataegus monogyna</i>), Hazel (<i>Corylus avellana</i>), Blackthorn (<i>Prunus spinosa</i>), Dogwood (<i>Cornus sanguinea</i>)	3	Multi	0.1	1	1	1	1	0	Young	Fair	Short section of field boundary hedgerow.	No action required at this time.	20-40	C
T16	Field Maple (<i>Acer campestre</i>)	14	Multi	0.6	5	5	5	5	2	Mature	Good	Prominent field boundary hedgerow exhibiting some minor animal damage on main stem.	No action required at this time.	>40	B
G17	Group of: Hazel (<i>Corylus avellana</i>), Hawthorn (<i>Crataegus monogyna</i>), Dogwood (<i>Cornus sanguinea</i>), Goat Willow (<i>Salix caprea</i>)	9	Single and Multi	0.2 (avg.)	3	2	3	2	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	20-40	C

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T18	Sycamore (<i>Acer pseudoplatanus</i>)	18	Single	0.64	9	7	8	3	3	Mature	Fair	Field boundary tree of reasonable form.	No action required at this time.	>40	B
T19	Sycamore (<i>Acer pseudoplatanus</i>)	18	Multi	0.75	3	3	8	8	4	Middle aged	Fair	Prominent twin-stemmed field boundary tree of reasonable form.	No action required at this time.	>40	B
T20	Oak (<i>Quercus robur</i>)	8	Single	0.53	3	1	4	5	3	Middle aged	Fair to poor	Scrubby field boundary specimen that has been severely pruned in the past.	Monitor for stability.	10-20	C
G21	Group of: Hawthorn (<i>Crataegus monogyna</i>), Hazel (<i>Corylus avellana</i>), Holly (<i>Ilex aquifolium</i>)	5	Multi	0.15	1	1	1	1	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	20-40	C
T22	Sycamore (<i>Acer pseudoplatanus</i>)	16	Single	0.32	4	5	4	4	6	Middle aged	Fair	Field boundary tree of reasonable form.	No action required at this time.	20-40	B
G23	Group of: Sycamore (<i>Acer pseudoplatanus</i>)	16	Single and Multi	0.4 (avg.)	10	2	2	5	2	Middle aged	Poor	Trees of poor form exhibiting significant dieback in upper crowns and splits within main stems. These specimens are unsafe for long-term retention.	Remove.	<10	U
T24	Dead														
G25	Group of: Sycamore (<i>Acer pseudoplatanus</i>), Holly (<i>Ilex aquifolium</i>)	Up to 12	Single and Multi	0.2 (avg.)	3	2	2	3	1	Middle aged	Fair to poor	Scrubby specimens forming short section of overgrown hedgerow. Sycamores exhibit some squirrel damage within crowns.	Monitor for health.	10-20	C

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G26	Group of: Dogwood (<i>Cornus sanguinea</i>), Hazel (<i>Corylus avellana</i>)	2	Multi	0.1	1	1	1	1	0	Young	Fair	Short section of young hedgerow.	No action required at this time.	20-40	C
G27	Group of: Sycamore (<i>Acer pseudoplatanus</i>), Hawthorn (<i>Crataegus monogyna</i>), Ash (<i>Fraxinus excelsior</i>)	4	Single and Multi	0.1	1	1	1	1	0	Middle aged	Fair	Scrubby specimen forming short section of gappy hedgerow.	Monitor health of Ash.	20-40	C
T28	Ash (<i>Fraxinus excelsior</i>)	16	Multi	0.45	5	5	4	3	2	Middle aged	Poor	Field boundary tree exhibiting early stage symptoms of Ash Dieback Disease.	Remove.	<10	U

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G29	Group of: Elder (<i>Sambucus nigra</i>), Hazel (<i>Corylus avellana</i>), Beech (<i>Fagus sylvatica</i>), Sycamore (<i>Acer pseudoplatanus</i>), Elm (<i>Ulmus</i> spp.), Hawthorn (<i>Crataegus monogyna</i>), Blackthorn (<i>Prunus spinosa</i>), Goat Willow (<i>Salix caprea</i>)	3	Multi	0.15	1	1	1	1	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	20-40	C
T30	Sycamore (<i>Acer pseudoplatanus</i>)	17	Single	0.48	6	8	6	3	4	Middle aged	Fair	Field boundary tree of reasonable form and vigour.	No action required at this time.	>40	B
T31	Pear (<i>Pyrus</i> spp.)	9	Single	0.4 (est.)	2	2	1	1	3	Mature	Fair	Tree located within private residential garden thus preventing full inspection and accurate measurement. Fruit tree of variable form.	No action required at this time.	10-20	C
T32	Sycamore (<i>Acer pseudoplatanus</i>)	18	Single	0.6 (est.)	7	7	7	7	4	Mature	Fair	Tree located within private residential garden thus preventing full inspection and accurate measurement. Prominent specimen of reasonable form and vigour.	No action required at this time.	>40	B

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T33	Scots Pine (<i>Pinus sylvestris</i>)	8	Single	0.2 (est.)	2	2	2	2	2	Young	Fair	Tree located within private residential garden thus preventing full inspection and accurate measurement. Tree of reasonable form.	No action required at this time.	>40	B
T34	Sycamore (<i>Acer pseudoplatanus</i>)	17	Multi	0.6 (est.)	5	5	5	5	3	Middle aged	Fair	Tree located within private residential garden thus preventing full inspection and accurate measurement. Tree of reasonable form and vigour.	No action required at this time.	>40	B
T35	Japanese Maple (<i>Acer palmatum</i>)	3	Multi	0.15	1	1	1	1	1	Young	Fair	Ornamental tree located within private residential garden thus preventing full inspection and accurate measurement. Tree of reasonable form.	No action required at this time.	20-40	C
G36	Group of: Lawson Cypress (<i>Chamaecyparis lawsoniana</i>), Dogwood (<i>Cornus sanguinea</i>), Elder (<i>Sambucus nigra</i>), Hazel (<i>Corylus avellana</i>)	2	Single and Multi	0.1	1	1	1	1	0	Young	Fair	Short section of young hedgerow.	No action required at this time.	20-40	C
T37	Sycamore	15	Multi	0.55	5	4	4	6	3	Middle aged	Fair	Twin-stemmed hedgerow specimen that has been severely pruned on southern side due to presence of overhead cables.	Monitor for stability.	20-40	C

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G38	Group of: Hazel (<i>Corylus avellana</i>), Holly (<i>Ilex aquifolium</i>), Sycamore (<i>Acer pseudoplatanus</i>)	5	Multi	0.15	1	1	1	1	0	Middle aged	Fair	Short section of scrubby hedgerow.	Monitor for stability.	20-40	C
T39	Alder (<i>Alnus glutinosa</i>)	18	Multi	0.9	10	9	10	9	4	Mature	Fair	Tree located within private residential garden thus preventing full inspection and accurate measurement. Tree of reasonable form.	Monitor for stability.	20-40	B
G40	Group of: Hawthorn (<i>Crataegus monogyna</i>), Hazel (<i>Corylus avellana</i>)	5	Single and Multi	0.15	3	1	2	3	0	Middle aged	Fair	Short section of scrubby hedgerow located to the west of drainage ditch.	No action required at this time.	20-40	C
G41	Group of: Hazel (<i>Corylus avellana</i>), Hawthorn (<i>Crataegus monogyna</i>), Blackthorn (<i>Prunus spinosa</i>)	1	Multi	0.1	0.5	0.5	0.5	0.5	0	Middle aged	Fair	Section of gappy hedgerow located on residential boundary.	No action required at this time.	20-40	C

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T42	Larch (<i>Larix</i> spp.)	17	Single	0.5 (est.)	6	6	7	7	6	Mature	Fair to poor	Tree located within private residential garden thus preventing full inspection and accurate measurement. A tree of poor form that has previously been heavily reduced in upper crown.	Monitor for stability.	10-20	C
T43	Oak (<i>Quercus robur</i>)	20	Single	0.75 (est.)	9	9	10	10	2	Mature	Fair	Prominent specimen located on boundary of private residential garden thus preventing full inspection and accurate measurement. A tree of reasonable form. Some deadwood extending over access lane.	Prune to remove unstable deadwood of diameter >50mm extending over access lane.	>40	A
G44	Group of: Hawthorn (<i>Crataegus monogyna</i>)	6	Multi	0.15	1	1	1	1	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	20-40	C
G45	Group of: Hazel (<i>Corylus avellana</i>), Holly (<i>Ilex aquifolium</i>)	11	Multi	0.3	2	5	5	5	0	Middle aged	Fair	Scrubby specimens forming gappy overgrown hedgerow.	Monitor for stability.	20-40	C
G46	Group of: Sycamore (<i>Acer pseudoplatanus</i>)	20	Multi	0.7	7	8	8	9	2	Middle aged	Fair	Linear feature located on field boundary containing trees of generally reasonable form.	Monitor for stability.	20-40	B
T47	Sycamore (<i>Acer pseudoplatanus</i>)	20	Multi	0.65	9	8	9	6	3	Mature	Fair	Prominent twin-stemmed specimen located on private residential garden boundary thus preventing full inspection. Some squirrel damage to upper branches on northern side of crown.	Prune to remove unstable deadwood of diameter >50mm.	20-40	B

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G48	Group of: Sycamore (<i>Acer pseudoplatanus</i>)	21	Single and Multi	0.5 (avg.)	8	6	6	9	2	Middle aged	Fair	Linear feature located on field boundary containing trees of generally reasonable form.	No action required at this time.	>40	B
G49	Group of: Sycamore (<i>Acer pseudoplatanus</i>), Holly (<i>Ilex aquifolium</i>)	16	Single and Multi	0.3 (avg.)	4	4	4	4	3	Middle aged	Fair	Linear feature located either side of drainage ditch. Trees of generally reasonable form and vigour.	No action required at this time.	20-40	B
G50	Group of: Apple (<i>Malus</i> spp.), Cherry (<i>Prunus</i> spp.)	4	Single	0.2 (est.)	2	2	2	2	1	Middle aged	Fair	Trees located within private residential garden thus preventing full inspection and accurate measurement.	No action required at this time.	20-40	C
G51	Group of: Holly (<i>Ilex aquifolium</i>), Hazel (<i>Corylus avellana</i>), Sycamore (<i>Acer pseudoplatanus</i>)	2	Multi	0.15	1	1	1	1	0	Middle aged	Fair	Scrubby specimens forming gappy boundary hedgerow to private residential garden.	No action required at this time.	10-20	C
G52	Group of: Hawthorn (<i>Crataegus monogyna</i>)	4	Single and Multi	0.15	1	1	1	1	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	20-40	C
G53	Group of: Holly (<i>Ilex aquifolium</i>), Hazel (<i>Corylus avellana</i>), Hawthorn (<i>Crataegus monogyna</i>)	10	Single and Multi	0.2	2	2	2	2	0	Middle aged	Fair to poor	Scrubby specimens forming gappy hedgerow either side of watercourse. Some dieback within crowns.	Monitor for health.	10-20	C

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T54	Elm (<i>Ulmus</i> spp.)	16	Single	0.45 (est.)	5	5	5	5	4	Middle aged	Fair	Tree located within private residential garden thus preventing full inspection and accurate measurement. Tree of reasonable form. Some slight thinning of foliage within crown.	Monitor for health.	20-40	B
T55	Oak (<i>Quercus robur</i>)	16	Single	0.4 (est.)	3	5	5	5	5	Middle aged	Good	Tree located within private residential garden thus preventing full inspection and accurate measurement. Prominent specimen of reasonable form.	No action required at this time.	>40	B
T56	Holly (<i>Ilex aquifolium</i>)	10	Multi	0.55	3	3	5	5	1	Middle aged	Fair to poor	Scrubby specimen exhibiting significant thinning of foliage throughout crown.	Monitor for health.	10-20	C
G57	Group of: Holly (<i>Ilex aquifolium</i>), Hazel (<i>Corylus avellana</i>), Elm (<i>Ulmus</i> spp.), Hawthorn (<i>Crataegus monogyna</i>)	4	Single and Multi	0.15	1	1	1	1	0	Middle aged	Fair	Scrubby specimens forming gappy hedgerow.	No action required at this time.	20-40	C
G58	Group of: Sycamore (<i>Acer pseudoplatanus</i>), Oak (<i>Quercus robur</i>)	23	Single and Multi	0.9	8	11	11	11	4	Mature	Good	Closely spaced group of trees forming a single collective crown. Prominent specimens of reasonable form.	No action required at this time.	>40	A

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T59	Alder (<i>Alnus glutinosa</i>)	19	Single	0.47	6	5	5	6	3	Mature	Fair	Prominent specimen of reasonable form.	No action required at this time.	20-40	B
T60	Alder (<i>Alnus glutinosa</i>)	18	Multi	0.9	9	6	8	8	2	Mature	Fair	Prominent riverside tree of reasonable form.	No action required at this time.	20-40	B
G61	Group of: Sycamore (<i>Acer pseudoplatanus</i>)	17	Single and Multi	0.35	8	8	8	7	2	Middle aged	Fair to poor	Riverside specimens of variable form exhibiting some dieback within crowns.	Monitor for health and stability.	10-20	C
G62	Group of: Crack Willow (<i>Salix fragilis</i>)	13	Multi	0.45	5	5	5	5	2	Middle aged	Poor	Multi-stemmed specimens with weak basal forks that are vulnerable to structural failure. This species is unsuitable for long-term retention.	Remove.	<10	U
T63	Alder (<i>Alnus glutinosa</i>)	11	Multi	0.35	4	3	4	4	3	Middle aged	Fair to poor	Twin-stemmed specimen of variable form that has suffered fire damage at base.	Monitor for health.	10-20	C
T64	Sycamore (<i>Acer pseudoplatanus</i>)	20	Multi	0.8	9	9	9	9	3	Mature	Fair	Prominent riverside tree of reasonable form. Some fire damage in lower crown on eastern side which is not considered significant.	Monitor for health.	>40	B
G65	Group of: Alder (<i>Alnus glutinosa</i>), Ash (<i>Fraxinus excelsior</i>)	13	Single and Multi	0.15 (avg.)	2	2	2	2	2	Middle aged	Fair to poor	Linear feature dominated by Ash infected with Ash Dieback Disease.	Remove specimens of infected Ash trees.	10-20	C

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G66	Group of: Sycamore (<i>Acer pseudoplatanus</i>), Alder (<i>Alnus glutinosa</i>)	Up to 19	Single and Multi	0.6 (avg.)	6	6	6	6	3	Mature	Fair	Prominent group of trees sited adjacent to riverbank.	No action required at this time.	20-40	C
T67	Ash (<i>Fraxinus excelsior</i>)	17	Single	0.34	6	3	2	2	7	Middle aged	Poor	Tree of variable form exhibiting early stage symptoms of Ash Dieback Disease.	Remove.	<10	U
T68	Ash (<i>Fraxinus excelsior</i>)	9	Single	0.14	1	3	1	1	3	Young	Poor	Heavily suppressed specimen exhibiting early stage symptoms of Ash Dieback Disease.	Remove.	<10	U
T69	Dead														
T70	Ash (<i>Fraxinus excelsior</i>)	16	Single	0.36	2	7	7	3	3	Middle aged	Poor	Tree of variable form exhibiting symptoms of Ash Dieback Disease.	Remove.	<10	U
G71	Group of: Alder (<i>Alnus glutinosa</i>), Sycamore (<i>Acer pseudoplatanus</i>)	18	Single and Multi	0.45 (avg.)	6	6	6	6	2	Middle aged	Fair	Closely spaced trees forming a unified crown.	No action required at this time.	>40	B
T72	Dead														
T73	Ash (<i>Fraxinus excelsior</i>)	20	Multi	0.65	5	7	8	6	5	Mature	Poor	Prominent twin-stemmed specimen exhibiting early stage symptoms of Ash Dieback Disease.	Remove.	<10	U
G74	Group of: Holly (<i>Ilex aquifolium</i>)	16	Multi	0.35	4	4	4	4	2	Middle aged	Fair	Trees of reasonable form and vigour.	No action required at this time.	>40	B

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G75	Group of: Ash (<i>Fraxinus excelsior</i>), Elm (<i>Ulmus</i> spp.)	6	Single	0.1	4	2	0	0	2	Young	Poor	Ash is infected with Ash Dieback Disease. Elm exhibits significant basal decay.	Remove.	<10	U
G76	Group of: Sycamore (<i>Acer pseudoplatanus</i>)	19	Multi	0.65	8	8	8	8	3	Middle aged	Fair	Trees of reasonable form located on banks of watercourse. Dense vegetation at base prevents full inspection.	Monitor for stability.	20-40	B
T77	Ash (<i>Fraxinus excelsior</i>)	20	Single	0.58	7	6	7	6	7	Mature	Poor	Tree located adjacent to watercourse. This specimen is infected with Ash Dieback Disease.	Remove.	<10	U
T78	Oak (<i>Quercus robur</i>)	18	Single	0.73	7	8	8	10	2	Mature	Fair	Prominent specimen located adjacent to watercourse. Tree of reasonable form and vigour.	No action required at this time.	>40	B
T79	Field Maple (<i>Acer campestre</i>)	17	Single	0.43	5	5	5	5	2	Mature	Fair	Prominent specimen located at edge of watercourse.	No action required at this time.	>40	B
G80	Group of: Hazel (<i>Corylus avellana</i>), Hawthorn (<i>Crataegus monogyna</i>), Holly (<i>Ilex aquifolium</i>), Cherry (<i>Prunus</i> spp.)	10	Single and Multi	0.2 (avg.)	3	3	3	3	0	Middle aged	Fair	Scrubby specimens forming gappy hedgerow located either side of watercourse.	No action required at this time.	20-40	C

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G81	Group of: Blackthorn (<i>Prunus spinosa</i>)	4	Multi	0.15	1	1	1	1	1	Middle aged	Fair to poor	Scrubby specimens that have suffered some animal damage in lower crowns.	Monitor for health.	10-20	C
T82	Alder (<i>Alnus glutinosa</i>)	7	Multi	0.2	2	2	2	2	2	Young	Fair to poor	Self-sown twin-stemmed specimen of variable form.	No action required at this time.	10-20	C
T83	Sycamore (<i>Acer pseudoplatanus</i>)	11	Multi	0.3	4	4	4	4	1	Middle aged	Fair	Scrubby isolated specimen of variable form.	No action required at this time.	10-20	C
T84	Crack Willow (<i>Salix fragilis</i>)	10	Multi	0.2	1	1	1	1	0	Young	Fair to poor	Multi-stemmed specimen located immediately adjacent to riverbank. This species is vulnerable to structural failure as it matures.	Monitor for stability.	10-20	C
T85	Goat Willow (<i>Salix caprea</i>)	8	Multi	0.3	2	2	2	2	0	Middle aged	Fair to poor	Tree of variable form located immediately adjacent to riverbank. This species is vulnerable to structural failure as it matures.	Monitor for stability.	10-20	C
T86	Ash (<i>Fraxinus excelsior</i>)	19	Multi	0.65	3	7	8	7	3	Mature	Poor	Woodland edge tree severely infected with Ash Dieback Disease.	Remove.	<10	U
G87	Group of: Ash (<i>Fraxinus excelsior</i>)	22	Single	0.45	6	6	6	6	3	Middle aged	Poor	Woodland trees exhibiting symptoms of Ash Dieback Disease.	Remove.	<10	U

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G88	Group of: Alder (<i>Alnus glutinosa</i>), Sycamore (<i>Acer pseudoplatanus</i>), Hawthorn (<i>Crataegus monogyna</i>), Hazel (<i>Corylus avellana</i>)	Up to 20	Single and Multi	0.55 (avg.)	6	6	6	6	5	Mature	Fair	Woodland area constituted by trees of generally reasonable form and vigour.	No action required at this time.	>40	B
T89	Sycamore (<i>Acer pseudoplatanus</i>)	19	Multi	0.9	9	8	10	5	1	Mature	Good	Prominent specimen of reasonable form.	No action required at this time.	>40	B
T90	Oak (<i>Quercus robur</i>)	19	Single	1.07	10	11	11	11	3	Mature	Good	Prominent specimen of good form and well-balanced crown. Tree of good vigour.	No action required at this time.	>40	A
G91	Group of: Beech (<i>Fagus sylvatica</i>)	17	Single and Multi	0.45 (avg.)	5	5	5	5	2	Middle aged	Fair	Prominent linear feature formed by trees of generally reasonable form. Some evidence of slight defoliation within some specimens.	Monitor for stability.	20-40	B
T92	Lombardy Poplar (<i>Populus nigra</i> 'Italica')	24	Single	0.82	3	3	3	3	2	Mature	Fair	Prominent specimen of a species that is vulnerable to structural failure as it matures.	Monitor for safety.	10-20	C
T93	Lombardy Poplar (<i>Populus nigra</i> 'Italica')	24	Single	0.98	3	3	3	3	2	Mature	Fair	Prominent specimen of reasonable form. This species is vulnerable to structural failure as it matures.	Monitor for safety.	10-20	C
T94	Ash (<i>Fraxinus excelsior</i>)	17	Multi	0.45	8	5	6	6	3	Middle aged	Poor	Hedgerow tree exhibiting symptoms of Ash Dieback Disease.	Remove.	<10	U

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G95	Group of: Sycamore (<i>Acer pseudoplatanus</i>), Hawthorn (<i>Crataegus monogyna</i>), Hazel (<i>Corylus avellana</i>), Holly (<i>Ilex aquifolium</i>), Ash (<i>Fraxinus excelsior</i>)	10 (avg.)	Single and Multi	0.3	4	4	4	4	0	Middle aged	Fair	Trees and shrubs of reasonable form creating robust field boundary hedgerow.	No action required at this time.	>40	B
T96	Oak (<i>Quercus robur</i>)	23	Single	1.22	10	9	12	12	4	Mature	Good	Prominent specimen of good form and well-balanced crown.	No action required at this time.	>40	A
G97	Group of: Hawthorn (<i>Crataegus monogyna</i>), Hazel (<i>Corylus avellana</i>), Holly (<i>Ilex aquifolium</i>), Ash (<i>Fraxinus excelsior</i>), Sycamore (<i>Acer pseudoplatanus</i>)	Up to 12	Single and Multi	0.3 (avg.)	3	3	3	3	0	Middle aged	Fair	Trees of generally reasonable form creating robust field boundary hedgerow. Specimen of Ash exhibits early stage symptoms of Ash Dieback Disease.	Remove infected Ash tree.	>40	B
T98	Oak (<i>Quercus robur</i>)	14	Single	0.42	1	8	8	3	2	Middle aged	Fair	Hedgerow tree of variable form with notable sweep in main stem at 2m to the southeast.	Monitor for stability.	20-40	B

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T99	Oak (<i>Quercus robur</i>)	15	Single	0.59	8	5	8	8	3	Mature	Good	Prominent hedgerow tree of good form.	No action required at this time.	>40	A
G100	Group of: Goat Willow (<i>Salix caprea</i>)	9	Multi	0.25	3	3	3	3	1	Middle aged	Fair to poor	Scrubby specimens, some of which have partially collapsed.	Remove collapsed specimens. Monitor remaining trees for stability.	10-20	C
G101	Group of: Hazel (<i>Corylus avellana</i>), Goat Willow (<i>Salix caprea</i>), Hawthorn (<i>Crataegus monogyna</i>)	7	Multi	0.2	2	2	2	2	0	Middle aged	Fair	Scrubby specimens forming sporadic field boundary hedgerow.	No action required at this time.	20-40	C
T102	Oak (<i>Quercus robur</i>)	7	Multi	0.35	3	3	3	3	2	Young	Fair	Isolated specimen of reasonable form.	No action required at this time.	20-40	C
T103	Crack Willow (<i>Salix fragilis</i>)	18	Single	0.56	6	9	8	8	1	Mature	Poor	This specimen has partially collapsed in the past and is vulnerable to further structural failure.	Remove.	<10	U
G104	Group of: Hazel (<i>Corylus avellana</i>), Sycamore (<i>Acer pseudoplatanus</i>)	8	Single and Multi	0.2	1	1	1	1	1	Middle aged	Poor	Trees of poor form. Sycamores have suffered severe squirrel damage. Hazel is dead.	Remove.	<10	U
G105	Group of: Sycamore (<i>Acer pseudoplatanus</i>), Hazel (<i>Corylus avellana</i>)	10	Single and Multi	0.25	4	4	4	4	0	Middle aged	Fair to poor	Scrubby woodland edge specimens of variable form. Sycamores have suffered significant squirrel damage.	Monitor for safety.	10-20	C

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T106	Ash (<i>Fraxinus excelsior</i>)	19	Single	0.62	7	7	7	7	3	Mature	Poor	Prominent woodland edge specimen exhibiting early stage symptoms of Ash Dieback Disease.	Remove.	<10	U
G107	Group of: Crack Willow (<i>Salix fragilis</i>)	11	Single and Multi	0.3	7	3	2	3	2	Middle aged	Poor	Self-sown specimens of variable form vulnerable to structural failure.	Remove.	<10	U
G108	Group of: Crack Willow (<i>Salix fragilis</i>)	6	Multi	0.25	4	2	1	2	1	Middle aged	Poor	These specimens are partially collapsed.	Remove.	<10	U
T109	Ash (<i>Fraxinus excelsior</i>)	19	Single	0.72	9	6	6	9	4	Mature	Poor	Prominent field boundary tree severely infected with Ash Dieback Disease.	Remove.	<10	U
T110	Ash (<i>Fraxinus excelsior</i>)	17	Single	0.43	6	6	6	6	3	Middle aged	Poor	Prominent specimen exhibiting early stage symptoms of Ash Dieback Disease.	Remove.	<10	U
G111	Group of: Sycamore (<i>Acer pseudoplatanus</i>), Alder (<i>Alnus glutinosa</i>), Ash (<i>Fraxinus excelsior</i>), Hazel (<i>Corylus avellana</i>), Hawthorn (<i>Crataegus monogyna</i>)	12	Single and Multi	0.3	4	4	4	4	2	Middle aged	Fair	Trees of reasonable form creating robust overgrown field boundary hedgerow.	No action required at this time.	>40	B
T112	Alder (<i>Alnus glutinosa</i>)	14	Single	0.78	9	7	6	9	2	Mature	Fair	Prominent field boundary specimen of reasonable form.	No action required at this time.	20-40	B

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
T113	Ash (<i>Fraxinus excelsior</i>)	19	Multi	0.8	8	6	9	9	5	Mature	Poor	Prominent hedgerow tree severely infected with Ash Dieback Disease.	Remove.	<10	U
T114	Ash (<i>Fraxinus excelsior</i>)	19	Single	0.58	10	9	10	6	3	Mature	Poor	Prominent hedgerow tree severely infected with Ash Dieback Disease.	Remove.	<10	U
G115	Group of: Sycamore (<i>Acer pseudoplatanus</i>)	20	Single	0.65	8	7	10	9	3	Mature	Good	Trees forming unified crown. Notable specimens of reasonable form.	No action required at this time.	>40	A
G116	Group of: Oak (<i>Quercus robur</i>)	19	Single and Multi	0.75	9	8	7	10	2	Mature	Good	Prominent woodland edge trees of reasonable form.	No action required at this time.	>40	A
G117	Group of: Sycamore (<i>Acer pseudoplatanus</i>), Holly (<i>Ilex aquifolium</i>), Hazel (<i>Corylus avellana</i>)	10	Single and Multi	0.25	3	1	3	5	0	Middle aged	Fair	Scrubby woodland edge specimens leaning to the west.	No action required at this time.	10-20	C
T118	Oak (<i>Quercus robur</i>)	10	Single	0.46	4	4	4	6	2	Middle aged	Fair	Woodland edge tree of slightly squat habit.	No action required at this time.	>40	B
T119	Oak (<i>Quercus robur</i>)	19	Multi	0.95	10	10	10	12	2	Mature	Good	Prominent field boundary tree of reasonable form.	No action required at this time.	>40	A

Tree No.	Species	Height(m)	Single/Multi Stemmed	Stem Diameter(m)	Branch Spread(m)				Height of Crown(m)	Age	Physiological Condition	Structural Condition	Prel. Man. Recommendations	Est. Remaining Contribution	Category
					N	E	S	W							
G120	Group of: Holly (<i>Ilex aquifolium</i>), Hawthorn (<i>Crataegus monogyna</i>), Hazel (<i>Corylus avellana</i>)	9	Single and Multi	0.25	2	2	2	2	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	20-40	B
T121	Sycamore (<i>Acer pseudoplatanus</i>)	19	Multi	0.65	8	8	6	4	4	Mature	Poor	Prominent field boundary specimen exhibiting significant basal decay on southwestern side of main stem. This specimen is at risk of failure towards adjacent residential housing.	Remove.	<10	U
G122	Group of: Holly (<i>Ilex aquifolium</i>), Hazel (<i>Corylus avellana</i>), Hawthorn (<i>Crataegus monogyna</i>), Goat Willow (<i>Salix caprea</i>)	10	Single and Multi	0.3	3	3	3	3	0	Middle aged	Fair	Scrubby specimens forming gappy field boundary hedgerow.	No action required at this time.	20-40	C
G123	Group of: Hazel (<i>Corylus avellana</i>), Holly (<i>Ilex aquifolium</i>)	11	Multi	0.35	3	8	5	4	0	Mature	Fair to poor	Multi-stemmed specimens of variable form. Extensive thinning of foliage within Hollies. Easternmost Hazel leaning excessively to the east.	Remove excessively leaning stems. Monitor remaining stems for stability.	10-20	C